



7040 Virginia Manor Road  
Beltsville, MD 20705, USA  
Web: www.abpbio.com;  
Email: info@abpbio.com

## 5(6)-CR6G, SE [5-(and-6)-Carboxy-rhodamine 6G, succinimidyl ester]


Catalog Number	Packaging Size
C130	5 mg

**Storage upon receipt:** -20°C, protect from light

### Introduction

**5(6)-CR6G, SE** is a amine-reactive fluorescent probe for preparation of bioconjugates. The excitation and emission spectra of carboxyrhodamine 6G (CR 6G) fall between those of fluorescein and tetramethylrhodamine and with higher fluorescence quantum yield than tetramethylrhodamine conjugates. With a peak absorption at 525 nm, conjugated probes of carboxyrhodamine 6G are an excellent match to the 514 nm spectral line of the argon-ion laser.

### Specifications

<b>Label:</b>	Rhodamine 6G	
<b>Ex/Em:</b>	525/550 nm	
<b>Detection Method:</b>	Fluorescent	
<b>Solubility:</b>	DMSO, DMF	
<b>Molecular Formula:</b>	C <sub>31</sub> H <sub>29</sub> N <sub>3</sub> O <sub>7</sub>	
<b>Molecular Weight:</b>	555.59	
<b>CAS Number:</b>	349672-89-5	
<b>Storage Conditions:</b>	-20°C, protect from light	
<b>Shipping Condition:</b>	Room Temperature	

### Applications

Fluorescent labeling

### References:

1. Multiplex detection of single-nucleotide variations using molecular beacons.  
Marras SA, Kramer FR, Tyagi S  
Genet Anal (1999) 14:151-156
2. Intramolecular Excitonic Dimers in Protease Substrates: Modification of the Backbone Moiety to Probe the H-Dimer Structure.  
Packard BZ, et al.  
J Phys Chem B (1998) 102:1820-1820
3. Single-molecule detection and identification of multiple species by multiparameter fluorescence detection.  
Widengren J, Kudryavtsev V, Antonik M, Berger S, Gerken M, Seidel CA  
Anal Chem (2006) 78:2039-2050